

Drug Type / Street Name	Pharmaceutical Name	Detection Time (in urine)	Signs of Use
Amphetamin (AMP)			
Speed, Bennies, Dexies, Uppers, Footballs	Dexedrine, Benzedrine, Dextrostat, Adderall, Biphphetamine	2-4 days; cutoff level: 1000 ng/ml	Dilated pupils, decreased appetite; dry mouth & nose, bad breath; frequent lip licking, talkative, nervous; increase in heart rate & blood pressure
Barbiturates (BAR)			
Barbs, Downers, Tranqs, Blue Dolls, Bluebirds, rainbows, yellows, reds, red Devils, m&ms	Nembutal, luminal, Seconal, Tuinal, florinal, Neodorm, immenocital, Stadodorm	Short acting: 1 day (Seconal, Nembutal); long acting: 2-3 weeks (phenobarb); cutoff level: 300 ng/ml	pupils normal, but reaction to light slow; pulse, Bp and respiratory rate lowered, sedation, stupor, coma
Benzodiazepines (BZO)			
Bennies, rophies	librium, Valium, ativan, halicon, Vivol, Novopoxide, remestan, rohypnol, Xanax, Tranxene, restoril, Versed, clonopin, rivotril, Dalamane, Serax	Normal dose: 2-3 days; chronic use (1 or more yrs): 4-6 weeks; cutoff level: 300 ng/ml	Drowsiness, dizziness, muscle relaxation; hangover effect after large dose; rohypnol used as date rape drug
Buprenorphine (BUP)			
Buprenex, Temgesic	Suboxone, Subutex	1-6 days; 10 ng/ml	pinpoint pupils, extreme drowsiness, dizziness, blurred vision, slowed breathing
Cocaine (COC)			
coke, crack, Snow, flake, rock, Nose candy, powder, Blow	None	2-4 days; cutoff level: 300 ng/ml	Dilates pupils, increased heart rate and Bp; anesthesia, euphoria, sensitive to sound; and touch, less need for food, cardiotoxic
Gamma Hydroxybutyric Acid (GHB) *			
ghB, cherry meth, easy lay, Jib, georgia home Boy, Sleep 500, Soap, Vita g, grievous Bodily harm, Scoop, goop,	None	1 day	users feel euphoric, then sleepy; used as a date rape drug; slows breathing and heart rate

fantasy, everclear, Salty Water, liquid X			
Ketamine*			
Veterinary anesthetic, Kat, Special K, Vitamin K, K, horse Tranquilizer	Ketalar	up to 14 days; up to 30 days with chronic users; cutoff level: 25 ng/ml	analgesia, amnesia, hallucinations, foul- tasting drip from nasal passages; users feel they are outside of their body, vomiting, seizures, dissociation
Marijuana (THC)			
Weed, pot, grass, reefer, hash, mary Jane, cannabis, colombian, acapulco gold, ganja, Bhang	Marinol	1-2 joints: 2-3 days; oral ingestion: 1-5 days; moderate smoker: 5 days; heavy smoker(daily): 10 days; chronic smoker-5 joints per day: 14-18 days or as long as a month; cutoff level: 50 ng/ml	increased appetite, altered perception, memory loss, bloodshot eyes, giggle frequently, trouble walking, disorientation; increased heart rate, vasodilation, dry mouth and throat, decreased respiratory rate, panic attacks, and paranoia
Methadone (MTD)			
Amidone, Fizzes	Dolophine, methadone, l- polamidon, physeptone	2-3 days; cutoff level: 300 ng/ml	Similar to heroin but less intense; sedation analgesia, same as other opiates
Methamphetamine (MET)			
Ice, Crystal, Meth, Doe	Desoxyn, Methedrine	2-4 days; cutoff level: 1000 ng/mL	Same as amphetamines; overdoses cause elevated body temp.
Methylenedioxymethamphetamine (MDMA) (Ecstasy)			
XTC, E, Adam, Blue Lips, Blue Kisses, Disco Biscuits, X-ing, Doctor	None	1-3 days; cutoff level: 500 ng/mL	Same as amphetamines; may grind teeth and clench jaw, sweating and increase in body temp.
MDA*			
Eve, Love Pill	None	1-3 days	Same as amphetamines; hallucinations, increase in body temp.; tachycardia, agitation, very talkative
Opiates (OPI)			
(Heroin- and morphine-based drugs) Smack, H, Skag, Dope, Junk,	Hydromorphone (Dilaudid), Codeine, Tylenol with Codeine, Morphine (Avinza,	1-2 days; cutoff level: 2000 ng/mL or 300 ng/mL	Pupils constricted, pulse and BP lowered, body temp. lowered, euphoria initially and then drowsy, respiratory depression, convulsions, constipation, coma

Horse	Kadian, MS Contin, Oramorph SR, Hydrocodone (Vocodin): Pos. at 300 cutoff		
Oxycodone (OXY)			
OXY, OC ,OX, Oxycotton, Hillbilly Heroin, Killer	OxyContin, Percodan, Percocet, Roxicodone, Tylox, OxyContin is a time-released form of Oxycodone	1-3 days; OxyContin longer detection times; cutoff level: 100 ng/mL	Same as opiates
Phencyclidine (PCP)			
Yellow Footballs	Darvon, Darvocet	3-4 days; cutoff level: 300 ng/mL	Same as opiates
Tricyclic Antidepressants (TCA)			
	Amitriptyline (Elavil, Endep), Cyclobenzaprine (Flexeril, Clomipramine (Anafranil), Perphenazine (Triafon), Desipramine (Norpramin), Imipramine (Totranil), Nortriptyline (Pamelor), Protriptyline (Vivactil), Trimipramine (Surmontil), Promazine (Sparine), Nordoxepine, Doxepin (Sinequan)	Therapeutic dose: 3 days; Extended use: 4 weeks; cutoff: 1000 ng/mL	Elevated body temperature, elevated pulse and BP, pupils dilated, agitation, delirium, seizures, constipation, dry/ flush skin